


MYCOTOXINS IN SWINE DIETS

Why it is important now

The cool, wet weather we experienced this summer has caused many problems for the farmers in this area. This fall there is one more – there are reports of grain that is contaminated with fusarium mould.



Deoxynivalenol (a.k.a. DON, vomitoxin) is a mycotoxin that can be produced when fusarium moulds contaminate cereal grains, including wheat, barley, and corn. Pigs are more sensitive to DON contamination in their feed than other farm animals. Growing beef cattle, sheep, and poultry have an Agriculture Canada Guideline of 5 ppm, while the guideline for pigs is 1 ppm (1 mg/kg).

The most common symptom pigs show when given DON contaminated feed is a reduction of feed intake and a corresponding decrease in weight gain. Some decrease in feed intake will likely be seen if DON contamination exceeds 1 ppm. In research conducted at the Prairie Swine Centre, feed intake and daily gain of late-nursery pigs decreased 9.1 and 5.2%, respectively, when pigs were fed diets containing 1.57 ppm DON for 22 days. Although pigs may vomit at high levels of DON contamination (~20 ppm) it is more likely that they will refuse feed completely (~12 ppm) before that occurs. Younger animals will be more severely affected than older animals. While the general recommendation for swine is to limit DON in diets for pigs to less than 1 ppm, a maximum of 0.5 is preferable for nursery pigs. While there doesn't seem to be direct negative reproductive effects from feeding DON contaminated diets to breeding stock, the reduced feed intake itself may be a problem. Therefore, DON contaminated feeds should be avoided in diets for breeding stock whenever possible.



Cross contamination

Often when feed is contaminated with one type of mycotoxin another type will also be present. For example, feed contaminated with DON can also be contaminated with zearalenone, another mycotoxin caused by fusarium.

As with DON, zearalenone affects pigs more than other farm animals. Zearalenone has estrogenic effects and when present in the feed causes vulva enlargement in pre-pubescent gilts, as well as decreased litter size and infertility. Keep zearalenone concentrations under 0.5 ppm for all swine and avoid using zearalenone contaminated grains altogether in diets for breeding and replacement swine.

Sampling

It is difficult to get a good sample of grain to test for mycotoxin contamination. There is usually a great deal of variation in the amount of mycotoxin present from one area to another within a truck or bin. Therefore, it is easy to get false negatives or lower or higher values than is actually present in the overall grain. Combining small samples from several different areas will help you be sure that a representative sample has been obtained.

Funding Provided by:



The following (Table 1) is an example of our sampling data from 10 different 1-tonne totes of corn obtained from a single source. The differences between labs may be partially explained by differences in procedures and timing of analysis.

Table 1. DON concentrations (ppm) in samples of contaminated corn.

Sample #	Lab #1	Lab #2	% Difference
1	5.5	6.6	18
2	4.1	7.3	56
3	2.4	6.3	90
4	4.6	8.1	55
5	4.2	2.2	63
6	4.8	7.7	46
7	4.5	9.6	72
8	5.3	7.7	37
9	4.6	6.9	40
10	4.8	6.6	32
Mean	4.48	6.90	51
Coefficient of Variation	18.96%	27.67%	

What to do

There are no feed additives approved in Canada to decrease the impact of DON on swine. Cleaning grain to decrease dust and small shrivelled kernels helps lower the concentration of DON in the grain. If possible, feed contaminated grains to less sensitive species and use clean grains in swine diets.



References:

van Heugten, E., 2000. Mycotoxins and other antinutritional factors in swine feeds. Pages 563 – 583. *in*: Swine Nutrition. A. J. Lewis, L. L. Southern, ed. CRC Press, Boca Raton, FL

Dupchak, K., Feeding Fusarium-Contaminated Grain to Livestock.

<http://www.gov.mb.ca/agriculture/livestock/nutrition/bza00s01.html> accessed 29/09/2010

CAST. Mycotoxins – risks in plants, animal, and humans. 2003. Ames, Iowa, USA.

Fast Facts

- Mould produces mycotoxins but high mould content doesn't mean mycotoxins are present and mycotoxins can be present when there is no longer any mould.
- Not all moulds produce mycotoxins.
- Swine are the farm animal most sensitive to DON (a.k.a. Deoxynivalenol, vomitoxin).
- Limit DON to less than 1 ppm in diets for swine.

Knowledge just a click away



Turn knowledge into a powerful tool to help you improve your bottom line.

PRAIRIE SWINE CENTRE

Prairie Swine Centre
Phone: (306) 373-9922
Fax: (306) 955-2510

prairieswine.ca

Pork Insight brings you the world's knowledge

The Prairie Swine Centre is dedicated to supporting a profitable Canadian pork industry with reliable information to reduce costs and improve productivity. Check out the new Pork Insight research website, powered by prairieswine.ca. If you're looking for practical, science-based information, the Pork Insight database contains over 5,500 articles. It's comprehensive, easy to search, and will help improve your operation's bottom line.

Pork Insight www.prairieswine.ca



Agriculture and Agri-Food Canada



AGRICULTURE COUNCIL OF SASKATCHEWAN INC.



Canada



ALMA
Alberta Livestock and Meat Agency Ltd.