

FEEDING MYCOTOXIN CONTAMINATED GRAIN



Mycotoxins are chemicals (secondary metabolites) produced by moulds or fungi infecting grains. There are over 400 known mycotoxins; however only a small number of these probably affect pig performance on a regular basis. It is important to note that the presence of the mould or fungi does not guarantee the presence of mycotoxins; conversely, mycotoxins can be present in a sample with no obvious mould contamination. The mycotoxins of major concern in Canada are listed in Table 1. Several factors contribute to the production of mycotoxins in grain, including humidity and temperature during the growing and harvest periods, oxygen availability during growth, harvest, transport or storage and insect or bird damage. Multiple mycotoxins may be present at the same time and mycotoxins may be “masked”. These are mycotoxins bound to another molecule which may make them undetectable by routine assays. They will however, break down in the gut, and cause problems.

Pigs are more susceptible to the effects of most mycotoxins than other species, especially ruminants. The age of the animal and production status are important considerations. Table 2 outlines the mycotoxins of major concern, and their impact on animal performance.

In order to determine optimal feeding strategies, it is critical to know which mycotoxins are present and the approximate concentration. Many commercial laboratories can analyze for the common mycotoxins. The difficulty is obtaining a sample which is representative of the entire lot. When sampling grains or feeds, subsamples from 12-20 locations should be collected, composited and mixed thoroughly (Whitlow et al., 2014). Once a sample is collected, it is also important to store it in a dry, cool area to impede further mycotoxin development before the analysis. Mycotoxins are often distributed unevenly throughout the load, and very small quantities can cause problems (1 part per million (ppm) is equal to 1 contaminated grain in 1 million non-contaminated grains). The more subsamples collected, the better the likelihood of obtaining a laboratory analysis which really represents what is in the feed.

The CFIA has regulatory guidelines for the feeding of mycotoxins to livestock. This document reminds us that mycotoxin contamination is typically higher in the lighter fractions (grain dust, screenings, shrivelled kernels, etc.), and that while removing these fractions from the parent stock may help to reduce overall

Mycotoxins, which are produced from moulds, can contaminate all grains and grain-by-products commonly fed to swine in Western Canada. Personnel working with grains should avoid inhaling the dust and wear a mask. Dilution is (only partially) the solution.

contamination, it also means that these fractions are typically heavily contaminated. Because mycotoxins and mould spores can concentrate in grain dust it is very important that inhalation is avoided and dust masks are worn when handling, as they will affect human health also. Soaking, dehulling, cleaning and/or roasting may be beneficial in some cases, as are some dietary additives.

The Canadian regulatory guidelines for feeding mycotoxins are summarized in Table 1. Diets must not contain more than what is listed in these guidelines. If mycotoxin contamination is suspected; dilution can mitigate the problem, but because of the issues discussed with sampling, even when diluted, the grain should be fed to the least susceptible group (for example, keep ergot and ZEN out of your breeding herd).

Table 1. Legislated maximums, regulatory guidelines and recommended maximums for different mycotoxins into swine diets (adapted from Charmley and Trenholm, 2012)*

Mycotoxin	Commodity	Levels
Deoxynivalenol ¹	Diets for swine	1 ppm
Aflatoxins ²	Animal feeding stuffs	20 ppb
T-2 toxin ³	Swine diets	< 1 ppm
Zearalenone ³	Gilt diets	< 1-3 ppm
	Swine diets	< 0.25-5 ppm
Ochratoxin A ³	Swine diets (kidney damage)	0.2 ppm
	Swine diets (reduced weight gain)	2 ppm
Ergot Alkaloids ³	Swine diets	4-6 ppm
Fumonisin ³	Swine diets	10 ppm

*ppm is parts per million (mg/kg) and ppb is parts per billion.
¹Regulatory guidelines (Worldwide regulations for mycotoxins. FAO Food and Nutrition Paper 64, 1997)
²Legislated maximum tolerated level (Worldwide regulations for mycotoxins. FAO Food and Nutrition Paper 64, 1997)
³Recommended tolerance levels in Canada and the United States

Table 2. Major Effects of Mycotoxins on Swine Performance

Mycotoxin	Primary Effect	Stage Affected	Clinical Signs
Deoxynivalenol (DON, vomitoxin)	Affects serotonin receptors and cytokine production	All stages (younger pigs may be more susceptible)	<ul style="list-style-type: none"> • Reduced ADFI and ADG^{1,2} • Vomiting² • Diarrhea (soft or watery feces)³ • Reduced immune function³ • Mild changes to kidney, thyroid, blood⁴
Aflatoxins	Mutagenic and carcinogenic	All stages	<ul style="list-style-type: none"> • Reduced ADFI and ADG⁵ • Reduced milk production⁵ • Lethargy⁶ • Ataxia (lack of coordination)⁶ • Rough hair coat⁶ • Hemorrhage⁶ • Fatty liver⁶
Zearalenone	Estrogenic	Pre-pubertal gilts, sows and pre-pubertal boars	<ul style="list-style-type: none"> • Swelling and reddening of the vulva⁷ • Vaginal and/or rectal prolapse⁷ • Anestrus⁸ • Reduced litter size⁸ • Fetal resorption⁸ • Implantation failure⁸ • Decreased libido and testosterone⁹ • Feminization⁹
Ochratoxin A	Disrupts phenylalanine (an amino acid) metabolism	All stages	<ul style="list-style-type: none"> • Kidney damage¹⁰ • Decreased ADFI and ADG¹¹ • Immunosuppression, increased risk of infection¹²
Fumonisin	Disrupts lipid metabolism	All stages, especially young pigs	<ul style="list-style-type: none"> • Pulmonary edema³ • Reduced immunity³ • Decreased ADFI and ADG¹³ • Shortness of breath³ • Weakness³ • Cyanosis (blue/purple colour of skin/membranes)³
T-2 and HT-2 Toxins	Inhibits protein synthesis	All stages	<ul style="list-style-type: none"> • Unthriftiness⁶ • Low ADFI and ADG⁶ • Reproductive failure⁶ • Gastric upset (diarrhea)⁶ • Cellular necrosis⁶ • Immunosuppression⁶
Ergot Alkaloids	Neurological	All stages, especially the reproductive herd	<ul style="list-style-type: none"> • Lameness¹⁴ • Gangrene¹⁴ • Decreased ADG¹⁴ • Abortion¹⁴ • Agalactia (absence of milk production)¹⁴ • Ataxia¹⁴

¹Decreased ADFI and feed refusals have been shown at levels as low as 0.5-1 ppm (Smith et al., 2005), ²> 2-5 ppm is for decreased ADFI and ADG, vomiting and complete feed refusal at > 20 ppm (Haschek et al., 2002), ³Pierce and Diaz, 2014, ⁴JECFA, 2001, ⁵Nibbelink, 1986, ⁶Whitlow et al., 2014, ⁷Friend et al., 1990, ⁸Smith et al., 2005, ⁹Osweiler, 1986, ¹⁰Kidney damage occurs at levels as low as 0.5 ppm (Lippold et al., 1992), ¹¹Performance is affected at levels of 2 ppm or greater (Lippold et al., 1992; Stoev et al., 2000), ¹²Can occur when levels > 2 ppm are fed for longer periods of time (Harvey et al., 1992), ¹³ADG reduced by 11% when 10 ppm fumonisin B1 was fed to starter pigs for 8 weeks (Rotter et al., 1996), ¹⁴Strickland et al., 2011

For more information, please contact
Prairie Swine Centre

Box 21057, 2105 8th Street East
Saskatoon, Saskatchewan CANADA
S7H 5N9
Phone: (306) 373-9922
Fax: (306) 955-2510
E-mail: lee.whittington@usask.ca
Website: www.prairieswine.com

Prairie Swine Centre is an affiliate of
 **UNIVERSITY OF SASKATCHEWAN**



Saskatchewan
Ministry of
Agriculture

ONTARIO PORK